

## Genome Sequence of the Emerging Pathogen *Aeromonas caviae*<sup>∇</sup>

Scott A. Beatson,<sup>1</sup> Maria das Graças de Luna,<sup>2</sup> Nathan L. Bachmann,<sup>1</sup> Nabil-Fareed Alikhan,<sup>1</sup>  
Kirstin R. Hanks,<sup>1</sup> Mitchell J. Sullivan,<sup>1</sup> Bryan A. Wee,<sup>1</sup> Angela C. Freitas-Almeida,<sup>2</sup>  
Paula A. dos Santos,<sup>2</sup> Janyne T. B. de Melo,<sup>2</sup> Derrick J. P. Squire,<sup>3</sup>  
Adam F. Cunningham,<sup>3</sup> J. Ross Fitzgerald,<sup>4</sup>  
and Ian R. Henderson<sup>3\*</sup>

School of Chemistry and Molecular Biosciences, University of Queensland, Brisbane, Australia<sup>1</sup>; Departamento de Microbiologia, Imunologia e Parasitologia, Faculdade de Ciências Médicas, Universidade do Estado do Rio de Janeiro, Rio de Janeiro-RJ, Brazil<sup>2</sup>; School of Immunity and Infection, University of Birmingham, Birmingham, United Kingdom<sup>3</sup>; and The Roslin Institute and Centre for Infectious Diseases, University of Edinburgh, Edinburgh, United Kingdom<sup>4</sup>

Received 5 November 2010/Accepted 13 December 2010

***Aeromonas caviae* is a Gram-negative, motile and rod-shaped facultative anaerobe that is increasingly being recognized as a cause of diarrhea in children. Here we present the first genome sequence of an *A. caviae* strain that was isolated as the sole pathogen from a child with profuse diarrhea.**

*Aeromonads* are Gram-negative organisms belonging to the *Gammaproteobacteria*. Nineteen species of *Aeromonas* can be distinguished by biochemical and molecular techniques (5); most do not have a strong association with disease. However, *Aeromonas salmonicida* is a well-established fish pathogen, and *Aeromonas caviae*, *Aeromonas hydrophila*, and *Aeromonas veronii* account for 85% of isolates derived from human infections (5). While *Aeromonas* sp. can be isolated from ca. 2% of healthy individuals, association with stools from patients with diarrhea has been reported to be as high as 10.8% (4, 5). Here we report the genome sequence of *A. caviae* Ae398. This strain was isolated from a 17-month-old male at the Hospital Universitário Pedro Ernesto, Rio de Janeiro, Brazil. The patient presented with gastroenteritis and diarrhea, without mucus or blood, approximately three times a day for 3 weeks. *A. caviae* Ae398 was isolated as the sole pathogen from a stool specimen collected 3 weeks after onset of symptoms and was identified by conventional biochemical tests (3). The isolate presented extracellular enzymatic activities, resistance to several antibiotics, and adherence to epithelial cell monolayers (2, 3, 7, 9, 10).

Whole-genome shotgun sequencing of *A. caviae* Ae398 was performed by a combination of 454 GS-FLX (162,066 shotgun reads of average length 358 bp) and Illumina sequencing (25,272,910 51-bp paired-end reads), leading to a final assembly of 149 contigs >200 bp. The assembly represented 4,339,218 bp, with a mean contig size of 29,793 bp, an N50 value of 76,364 bp (where N50 is the contig length such that at least 50% of the bases of the assembly are contained within contigs of this size or greater), and an average G+C content of 61.4%. Nucleotide pairwise genome alignments (6) show the *A. caviae* Ae398 genome displays the highest overall synteny with

*A. hydrophila* ATCC 7966 (70.4% aligned) and *A. salmonicida* A449 (61.7% aligned) and average nucleotide identities of 88.8% and 86.8% across aligned regions, respectively.

The deduced size of the *A. caviae* Ae398 genome (4.43 Mb) is similar to those of *A. hydrophila* ATCC 7966 (4.47 Mb) and *A. salmonicida* A449 (4.70 Mb). *A. caviae* Ae398 harbors at least one conjugative plasmid of >30 kb, although it remains in several contigs in the assembly. Five different insertion (IS) element types are present in *A. caviae* Ae398, whereas *A. salmonicida* possesses 88 copies of 10 different IS elements and *A. hydrophila* completely lacks IS elements (8, 11). An ~33-kb putative prophage bounded by 55-bp repeats is found at the tRNA-Leu attachment site in *A. caviae* Ae398; the 3' region shares high identity with *phi*O18P (1), whereas the 5' region is most similar to structural genes from non-*Aeromonas* phage genomes.

*A. caviae* contains many putative virulence genes, including those encoding a type 2 secretion system (11), an RTX toxin, and polar flagella. The genome sequence of *A. caviae* Ae398 is an important milestone in understanding the diversity and pathogenesis of *A. caviae*, opening new avenues for investigating the pathogenic processes of this organism and allowing robust diagnostics to be developed.

**Nucleotide sequence accession numbers.** The assembly was deposited at the WGS division of DDBJ/EMBL/GenBank under accession no. CACP01000001 to CACP01000149.

This work was funded by the BBSRC and MRC grants to I.R.H. and by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ). S.A.B. is supported by an Australian Research Council Fellowship (DP0881347).

### REFERENCES

1. Beilstein, F., and B. Dreiseikelmann. 2008. Temperate bacteriophage *Phi*O18P from an *Aeromonas media* isolate: characterization and complete genome sequence. *Virology* **373**:25–29.
2. Castilho, M. C., et al. 2009. High frequency of hemolytic and cytotoxic activity in *Aeromonas* spp. isolated from clinical, food and environmental in Rio de Janeiro, Brazil. *Antonie Van Leeuwenhoek* **96**:53–61.
3. Freitas, A. C., S. M. Souza, L. C. Macedo, E. C. Pinto, and S. S. Pereira. 1998. *Aeromonas* species associated with gastroenteritis in children: prev-

\* Corresponding author. Mailing address: School of Immunity and Infection, University of Birmingham, Birmingham B15 2TT, United Kingdom. Phone: 44 1214724524. Fax: 44 1214143599. E-mail: i.r.henderson@bham.ac.uk.

<sup>∇</sup> Published ahead of print on 23 December 2010.

- alence, characteristics and virulence properties. *Rev. Microbiol.* **29**:152–157.
4. **Gracey, M., V. Burke, and J. Robinson.** 1982. *Aeromonas*-associated gastroenteritis. *Lancet* **ii**:1304–1306.
  5. **Janda, J. M., and S. L. Abbott.** 2010. The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clin. Microbiol. Rev.* **23**:35–73.
  6. **Kurtz, S., et al.** 2004. Versatile and open software for comparing large genomes. *Genome Biol.* **5**:R12.
  7. **Palu, A. P., et al.** 2006. Antimicrobial resistance in food and clinical *Aeromonas* isolates. *Food Microbiol.* **23**:504–509.
  8. **Reith, M. E., et al.** 2008. The genome of *Aeromonas salmonicida* subsp. *salmonicida* A449: insights into the evolution of a fish pathogen. *BMC Genomics* **9**:427.
  9. **Rocha-De-Souza, C. M., et al.** 2001. Identification of a 43-kDa outer-membrane protein as an adhesin in *Aeromonas caviae*. *J. Med. Microbiol.* **50**:313–319.
  10. **Rocha-De-Souza, C. M., et al.** 2003. Influence of polarisation and differentiation on interaction of 43-kDa outer-membrane protein of *Aeromonas caviae* with human enterocyte-like Caco-2 cell line. *Int. J. Mol. Med.* **11**:661–667.
  11. **Seshadri, R., et al.** 2006. Genome sequence of *Aeromonas hydrophila* ATCC 7966T: jack of all trades. *J. Bacteriol.* **188**:8272–8282.